

MSNL-18A

Residual Interactions in 4-Quasiparticle $K^T=14^-$ Isomer in $^{176}\text{Hf}^*$

ABSTRACT

We have for the first time located and identified a 4-quasiparticle isomer in a deformed nucleus. Its energy may be used as a test for the effective interaction between quasiparticles. We can account for the energy quite well with a simple δ -function interaction, but not with an interaction derived from Brueckner theory.

T. L. Khoo, F. M. Bernthal, and R. A. Warner
Cyclotron Laboratory and Departments of Physics and Chemistry
Michigan State University
East Lansing, Michigan 48824

and

G. P. Bertsch and G. Hamilton
Department of Physics and Cyclotron Laboratory
Michigan State University
East Lansing, Michigan 48824

*Work supported by U. S. National Science Foundation and U. S. Energy Research and Development Agency.

Information on the residual interaction in rare-earth deformed nuclei has come primarily from the singlet-triplet splittings of Gallagher-Moszkowski pairs in odd-odd nuclei,¹ and to a much lesser extent from similar splittings of two quasiparticle (qp) states in even-even nuclei.¹ The off-diagonal matrix elements of the residual interaction have also recently been obtained from the mixing between 2-neutron and 2-proton configurations^{2,4} in even-even nuclei. To date, however, no attempt has been made to extract information on the residual interactions in the rare-earth region from a study of states of seniority greater than two, despite the fact that many 3-qp states are known. To begin such an investigation it is, in fact, preferable to study a 4-qp system for which the energies of the constituent 2-qp states are already known. In this case, the effects of the residual interaction may be extracted from the observed 2- and 4-qp energies without having to calculate explicitly the individual qp energies.

The interactions thus deduced may then be used as a test of chosen effective interactions.

The twofold goals in the present investigation were to isolate experimentally a 4-qp state and, using its energy, to test the applicability of a 'realistic' force and a δ -force as effective interactions for rare-earth deformed nuclei. We prefer to search for very high K (≥ 12) states, since these should be rather pure due to the low density of such states even at the ~3 MeV excitation expected. An accompanying attractive feature is the fact that the large K should lead to isomerism, greatly simplifying the identification of the state.

The nucleus ^{176}Hf is a particularly favorable candidate for investigation since the lowest 2-quasiparticle excitations have been identified² and, furthermore, have high K so that one may expect the lowest 4-qp states to have very large K. In addition the decay of the 4-qp isomer should

proceed through known² bands built on the lower-lying 2-qp configurations, further simplifying the observation of the isomer.

The $^{176}\text{Yb}(\alpha, 4n)^{176}\text{Hf}$ reaction was used to populate high-K states in ^{176}Hf . The following experiments in which γ -rays were detected by Ge(Li) detectors were performed: a) observation of delayed γ -rays in a 5 msec interval between irradiation periods of 1 msec duration; b) search for short-lived isomers, in which the interval between beam bursts varied from 50 to 500 nsec; c) conventional 3-parameter $\gamma\gamma\gamma$ -coincidence; d) excitation function; and e) angular distribution measurements. In addition to two previously known³ 10- μ sec isomers, a new 401±6 μ sec isomer was discovered at 2866.0 keV. The decay scheme of the isomer is shown in Fig. 1. All the γ -rays in the figure which had sufficient intensity for half-life determination were found to decay with a 401- μ sec component. No other isomers with half-lives greater than a few nsec were found in ^{176}Hf . It is interesting to observe that the population of the 401- μ sec isomer accounts for an unusually large fraction (~30%) of the total ($\alpha, 4n$) cross-section.

The new isomer decays through two $K^{\pi}=8^-$ bands which had previously been identified² to spin 12. The energies, branching ratios and relative signs of the M1 and E2 matrix elements of the band members can all be accounted for in a model² in which there is increasing mixing between levels with increasing spin. On the basis of this model, there is little doubt that the 2827.3-keV level is the lower 13^- band member. (The higher 13^- level, not shown in Fig. 1, is found at 2920.9 keV.)

Using intensity balance considerations, the total internal conversion coefficient of the 38.7-keV transition was obtained ($\alpha_{\text{tot}}=12\pm 2$), leading to an unambiguous M1 assignment. From the M1 character of the

38.7-keV transition and the observed decay pattern, the most likely spin-parity for the 2866-keV isomer is 14^- . The 228- and 302-keV transitions should thus be L2 and their conversion coefficients ($c_{\text{tot}}(228) \approx 0.26$, $c_{\text{tot}}(302) \approx 0.16$) are consistent with this multipolarity. The isomerism of the 2866-keV level suggests K-forbiddenness, leading to a $K^{\pi}=14^-$ assignment. We have also identified⁵ the rotational band built on this isomer up to spin 19 and obtained an intrinsic g-factor, $g_K=0.59 \pm 0.02$, from intraband cascade-to-crossover ratios in the manner described in Ref. 2. This value is in excellent agreement with that expected for, and thus suggests, the 4-QP $K=14^-$ $\{7/2(404)p, 9/2(514)p, 7/2(514)n, 5/2(512)n\}$ configuration.

(It should be noted that no 2-QP states of such high K are expected at this energy.) This is the first time a high-K 4-QP isomer has been located in a deformed nucleus.⁶

The energy of a 4-QP state composed of 2 protons and 2 neutrons may be expressed⁷ in terms of the energies of the constituent 2-QP systems:

$$E(K_4) = E(K_{2p}) + E(K_{2n}) + [V(pn) - V_o(pn)] + \frac{1}{2}g(K_4 - K_{2p} - K_{2n}),$$

where $E(K_4)$ is the 4-QP energy, $E(K_{2p})$ and $E(K_{2n})$ are the observed energies of the 2-QP proton and neutron states, $[V(pn) - V_o(pn)]$ is the difference between the QP proton-neutron interaction energies in the ground and excited states, and the last term represents the correction for the zero-point rotation energy. The p-p and n-n interactions are already included in the 2-QP energies and need not be treated explicitly. The residual interactions are given by:

$$V(pn) - V_o(pn) = \sum_{i,k}^f [\langle ik | V | ik \rangle - \bar{\langle ik | 2v_i^2 v_k^2 \rangle}] + \sum_{i,k}^f V_{ik}$$

where $\langle ik | V | ik \rangle$ is the interaction matrix element for particles in orbits i and k, v_i^2 and v_k^2 are the pairing fullness parameters, and the bar denotes the time-reversed orbital. The prime indicates that the first sum contains only terms in which both orbitals are occupied in the 4-QP configuration; the second sum includes all remaining terms. Since many terms are involved in the second sum, we approximate individual matrix elements by an average one, $\langle ik | v | av \rangle$. In this approximation, it can be shown⁷ that

$$\sum_{i,k}^f V_{ik} = -\langle ik | v | av \rangle \sum_{i,k}^f (1-4v_i^2 v_k^2)$$

by noting that the number of interactions is equal in both the excited and ground states. The prime over the sum has the same meaning as in Eq. 1.

The residual interactions in a 4-QP state were evaluated using Eqs. 1 and 2 in the following manner. The single-particle Nilsson wave functions were expanded in terms of spherical states, from which the two-particle wave functions were derived in a j-j coupling scheme. The two-body matrix elements were evaluated using interactions between oscillator states of the relative coordinate derived from: a) a δ -interaction,

$$H(\xi) = 4\pi g(v/2)^{3/2} (1-\alpha \xi^2) \frac{\partial}{\partial \xi}$$

with the force strengths, $4\pi g(v/2)^{3/2} = 5.82$ MeV and $\alpha = 0.46$, adjusted to reproduce the splittings of singlet-triplet pairs in odd-odd¹ and even-even⁷ nuclei; and b) a 'realistic' interaction, including spin-orbit and tensor terms, obtained from the Reid potential using Brueckner theory in the manner described in Ref. 8. Details of the calculations will be published separately.⁷ The average interaction, $\langle ik | v | av \rangle$, was taken as the

mean of a large number of matrix elements calculated⁷ for 2-qp states of odd-odd nuclei. The required pairing factors in Eqs. 1 and 2 are taken from Ref. 4.

The energies of several 4-qp states in ^{176}Hf , in addition to that of the $K^{\pi}=14^-$ configuration of interest, have been calculated using Eqs. 1 and 2. These energies are presented in Table I, which also includes the predicted energy of a 4-qp $K^{\pi}=16^+$ state in ^{178}Hf . The energies, E_0 , derived from the sums of the constituent 2-qp energies without including residual p-n interactions are given in column 4 of the table. Columns 5 and 6 give the energies obtained after including p-n interactions calculated using the 'realistic' and the δ -function interactions, respectively. The use of the δ -force yields a calculated energy for the $K^{\pi}=14^-$ state in good agreement with experiment. Furthermore, whereas the sums of 2-qp energies predict the $K^{\pi}=12^+$ state to lie lowest, inclusion of the residual interactions makes the $K^{\pi}=14^-$ state lowest, in accord with observation.

(If the $K^{\pi}=12^+$ state, or any other high- K 4-qp state, were lowest in energy, it would have received significant feeding in the experiment and its isomeric decay via 2-qp bands would have been easily detectable.) The energies calculated using the 'realistic' interaction are very unsatisfactory, being too low by about 300 keV. Detailed analysis shows that this arises because the matrix elements of high- K p-n couplings calculated with the 'realistic' interaction tend to be too large compared with the average interaction, $\langle ik \rangle_{av}$, used in Eq. 2.

Similar calculations⁷ of the energies of some 3-qp states also show that the simple δ -force gives better agreement with experiment than the more sophisticated 'realistic' interaction. It would be of interest to determine the reasons for the deficiency of the realistic interaction in this mass region.

While the δ -function interaction is very successful in predicting 3- and 4-qp energies and also the splittings many Gallagher-Moszkowski pairs in odd-odd nuclei,^{1,7} it is unable to account^{1,9} for the odd-even shifts in $K=0$ bands of odd-odd nuclei. In fact, an interaction which can successfully account for the various types of data now available on deformed nuclei is yet to be found.

There is also a clear need for more data on other high- K 4-qp states and their constituent 2-qp components. A search for such configurations by means of (α, xn) reactions should be promising in the Hf region, where the orbitals near the Fermi level have large- Q projections. Preliminary analysis of on-going experiments⁵ indicate the existence of other high- K 4-qp structures in both ^{174}Hf and ^{176}Hf .

The authors would like to acknowledge helpful discussions with C. H. King.

REFERENCES

Table I. Low-Lying $K \geq 12$ Four-qp States in ^{176}Hf and ^{178}Hf

	K	Configuration ^a	E_o^b keV	E_{calc}^c keV	E_{calc}^d keV	E_{exp}^e keV
	^{176}Hf	14^-	$7/2_p$ $9/2_n$ $7/2_n$ $5/2_n$	3158	2580	2838
		14^-	$7/2_p$ $5/2_p$ $7/2_n$ $9/2_n$	3353	3105	3482
		12^+	$7/2_p$ $5/2_p$ $7/2_n$ $5/2_n$	3094	2560	2928
		15^+	$7/2_p$ $9/2_p$ $9/2_n$ $5/2_n$	3360	3012	3190
		16^+	$7/2_p$ $9/2_p$ $9/2_n$ $7/2_n$	3419	2697	3061
	^{178}Hf	16^+	$7/2_p$ $9/2_p$ $9/2_n$ $7/2_n$	2628	2069	2570
						2430-2550 ^f

^aThe single particle orbitals are $7/2_p$: $7/2(404)$, $9/2_p$: $9/2(514)$, $5/2_p$: $5/2(402)$, $7/2_n$: $7/2(514)$, $9/2_n$: $9/2(624)$, $5/2_n$: $5/2(512)$.

$$b E_o = E(K_{2p}) + E(K_{2n})$$

^cEvaluated using 'realistic' interaction to calculate matrix elements.

^dEvaluated using δ -interaction to calculate matrix elements.
^eEnergy > 2866 keV. Arguments are made in the text that the 2866-keV isomer is the lowest high- K 4-qp state.

^fRef. 3.

¹H. D. Jones, N. Onishi, T. Hess, and R. K. Sheline, Phys. Rev. C3, 529 (1971).

²T. L. Khoo, J. C. Waddington, R. A. O'Neil, Z. Prebrisz, D. G. Burke, and M. W. Johns, Phys. Rev. Lett. 28, 1717 (1972); T. L. Khoo, J. C. Waddington, and M. W. Johns, Can. J. Phys. 51, 2307 (1973).

³R. G. Helmer and C. W. Reich, Nucl. Phys. A114, 649 (1968).

⁴H. Massmann, J. O. Rasmussen, T. E. Ward, P. E. Haustein, and F. M. Bernthal, Phys. Rev. C9, 2312 (1974).

⁵T. L. Khoo, F. M. Bernthal, R. G. H. Robertson, and R. A. Warner, to be published.

⁶The decay of a $K^\pi = (16^+)$ isomer was previously observed (Ref. 3) in ^{178}Hf , but its exact location was not determined.

⁷T. L. Khoo, G. F. Bertsch, and G. Hamilton, to be published.

⁸G. F. Bertsch, The Practitioner's Shell Model, Amsterdam: North-Holland Publishing Co. (1972).

⁹N. D. Newby, Phys. Rev. 125, 2063 (1962).

FIGURE CAPTION

Figure 1 Decay scheme of 4qp $K^{\pi}=14^-$ isomer in ^{176}Hf . The γ -ray relative intensities of all transitions are indicated; the energies of the transitions directly de-exciting the isomer are given in parenthesis.

10




Fig. 1