

Formation of glutamine from  $^{13}\text{N}$ -labeled ammonia and dinitrogen, and from  $^{14}\text{C}$ -labeled glutamate, by heterocysts isolated from Anabaena cylindrica

[Running head: Formation of glutamine by heterocysts]

† Joseph Thomas, J. C. Meeks, C. Peter Volk\*, Paul W. Shaffer

MSU/ERDA Plant Research Laboratory

and

Sam M. Austin and W.-S. Chien

Cyclotron Laboratory and Department of Physics

Michigan State University, East Lansing

Michigan 48824

† Permanent Address: Biology and Agriculture Division  
Bhabha Atomic Research Centre  
Trombay, Bombay 400 085, India

A method is described for the isolation of metabolically active heterocysts from Anabaena cylindrica. These isolated heterocysts accounted for up to 34 percent of the acetylene reducing activity of whole filaments, and had a specific activity of up to 1560 nmoles  $\text{C}_2\text{H}_4$  formed  $(\text{mg heterocyst chlorophyll})^{-1} \cdot \text{min}^{-1}$ . Activity of glutamine synthetase is coupled to activity of nitrogenase in isolated heterocysts as shown by acetylene-inhibitable formation of  $^{13}\text{N}$  $\text{NH}_3$  and of amide-labeled  $^{13}\text{N}$ glutamine from  $^{13}\text{N}$  $\text{N}_2$ .

A method is also described for the production of 6-mCi amounts of  $^{13}\text{N}$  $\text{NH}_3$ . Isolated heterocysts formed  $^{13}\text{N}$ -labeled glutamine from  $^{13}\text{N}$  $\text{NH}_3$  and glutamate, and  $^{14}\text{C}$ -labeled glutamine from  $\text{NH}_3$  and  $^{14}\text{C}$ glutamate, in the presence of MgATP. Methionine sulfoximine strongly inhibited these syntheses.

Glutamate synthase is, after nitrogenase and glutamine synthetase, the third sequential enzyme involved in the assimilation of  $\text{N}_2$  by intact filaments. However, the kinetics of solubilization of the activity of glutamate synthase during cavitation of suspensions of A. cylindrica indicate that very little, if any, of the activity of that enzyme is located in heterocysts. Concordantly, isolated heterocysts failed to form substantial amounts of radioactive glutamate from either  $^{13}\text{N}$ glutamine or  $^{14}\text{C}$  $\alpha$ -ketoglutarate in the presence of other substrates and cofactors of the glutamate synthase reaction. However, they formed  $^{14}\text{C}$ glutamate rapidly from  $^{14}\text{C}$  $\alpha$ -

ketoglutarate by aminotransferase reactions, with various amino acids as the nitrogen donor. The implications of these findings with regard to the identities of the substances moving between heterocysts and vegetative cells are discussed.

Filamentous cyanobacteria which have been shown to be capable of fixing nitrogen gas ( $N_2$ ) aerobically are characterized by the presence of differentiated cells called heterocysts. A large amount of indirect evidence (6, 8, 22, 25, 27, 35) favors the hypothesis that heterocysts are major loci of nitrogen fixation in the filaments.

This hypothesis was substantiated by autoradiographic experiments with  $^{13}N$ -labeled  $N_2$  (32). Early attempts (7, 10, 20, 21) to demonstrate substantial nitrogenase activity in isolated heterocysts free of vegetative cells were unsuccessful. Recently, however, Peterson and Burris (16), working with a strain of Anabaena, described a procedure for isolating heterocysts which can reduce  $C_2H_2$ , an alternate substrate for nitrogenase, at up to 1840 nmoles  $[mg \text{ chlorophyll}(Chl)]^{-1} \cdot \text{min}^{-1}$ . When calculated on the basis of the heterocyst frequency in filaments, this rate accounted for 13 percent of the rate of  $C_2H_2$  reduction by intact filaments ( $1060 \text{ nmoles} \cdot \text{mg Chl}^{-1} \cdot \text{min}^{-1}$ ). Tel-Or and Stewart (23) found up to 45 percent of the activity of whole filaments in heterocysts isolated from Anabaena cylindrica, but the specific activity of nitrogenase in whole filaments reported by the latter workers was only  $0.194 \text{ nmoles } C_2H_4 \cdot (mg \text{ protein})^{-1} \cdot \text{min}^{-1}$ . This rate is approximately 6 nmoles  $C_2H_4 \cdot \text{mg Chl}^{-1} \cdot \text{min}^{-1}$ , if one assumes a ratio of protein to chlorophyll of approximately 30 (4).

The enzymatic pathway consisting of glutamine synthetase and glutamate synthase (glutamine amide:  $\alpha$ -ketoglutarate amido transferase) (24) was shown to be the major route of metabolism of ammonia in heterotrophic nitrogen-fixing bacteria (15). Using  $^{13}N$ -labeled nitrogen

gas ( $^{13}\text{N}_2$ ), we recently established that this pathway also mediates the metabolism of  $\text{N}_2$ -derived ammonia in *A. cylindrica* (26, 34). The presence of the two constituent enzymes in cyanobacteria had previously been demonstrated (3, 11).

In our experiments (34), ammonia derived from  $^{13}\text{N}_2$  was incorporated into glutamine so rapidly that the latter substance had as much  $^{13}\text{N}$  as had ammonia after about 2 s of fixation. In contrast, approximately 90 s elapsed before glutamate became as highly labeled as glutamine. These results suggested that glutamine synthetase is located close to the site of  $\text{N}_2$  fixation. The activity of glutamine synthetase in heterocysts had been found to be 1.7 times as great as that in vegetative cells, per mg protein (3). However, the localization of glutamate synthase in filamentous nitrogen fixing cyanobacteria has not heretofore been investigated.

In this paper we report that isolated heterocysts reduce  $\text{N}_2$  gas at a substantial fraction of intact-filament rates. Moreover, the nitrogenase activity in these heterocysts is metabolically coupled to activity of glutamine synthetase, so that the heterocysts produce  $^{13}\text{N}$ -labeled glutamine. However, the heterocysts produce very little or no glutamate, and concordantly have very low or no glutamate synthase activity. It was possible to obtain these results because we had developed a new method for isolating heterocysts of high metabolic activity, and which contained no more than 2 percent as many vegetative cells as heterocysts. Also important was development of a method for production of highly radioactive  $^{13}\text{N}$ -labeled ammonia.

#### MATERIALS AND METHODS

Cultures of *Anabaena cylindrica* Lemm. were grown aerobically in fermentors (for details, see 32) with  $\text{N}_2$  as nitrogen source, with an exponential doubling time of about 18 h, to a density of 0.5-0.76  $\mu\text{g Chl}\cdot\text{ml}^{-1}$ , except where noted otherwise. For experiments in which whole filaments were exposed to  $^{13}\text{N}_2$ , the cyanobacterium was concentrated to 27  $\mu\text{g Chl}\cdot\text{ml}^{-1}$  by centrifugation at 200 x g for 5 min, and was incubated aerobically, as described earlier (32), until used.

#### Isolation of heterocysts

Heterocysts were isolated as follows. The cyanobacterial suspension was sedimented at 200 x g for 25 min. The pellet was resuspended to 27-100  $\mu\text{g Chl}\cdot\text{ml}^{-1}$  in a portion of the supernatant fluid or in 5mM TES [N-tris(hydroxymethyl)-methyl-2-aminoethane sulfonic acid] buffer, pH 7.6. Ten ml of the concentrated suspension were thrice evacuated to 50  $\mu\text{m}$  of mercury, and regassed to 1 atm with argon. Further processing of the suspension was done under these strictly anaerobic conditions, with solutions and reagents degassed as described above. Biochemicals, unless otherwise specified, were obtained from Sigma Chemical Co., St. Louis, Mo. One ml of solution containing 2 or 5 mM TES, pH 7.6, and 200 mM sodium dithionite/was added to the 10-ml algal suspension, followed by 20 ml of a solution containing TES, D-mannitol (Fisher Scientific Co., Fair Lawn, N.J.) and  $\text{Na}_2\text{EDTA}$  [(ethylenedinitrilo)-tetraacetic acid, disodium salt]. The resulting suspension, containing 1.4 to 5 mM TES, 6.5 mM sodium dithionite, 26 mM  $\text{Na}_2\text{EDTA}$  and 650 mM D-mannitol, was incubated on a waterbath shaker (75 rpm) at 30°C. After 45 min, most of the vegetative cells were detached or

Such suspensions of filaments reduced acetylene at least as rapidly, per mg Chl, as when assayed immediately after withdrawal from a fermentor.

Reduction of acetylene by isolated heterocysts was normally measured anaerobically in the presence of dithionite, ATP, and an ATP generating system (see Table 1, footnote a). On occasion, certain of the components were omitted or replaced by others.

#### Labeling with $^{13}\text{N}_2$ and $^{13}\text{N}(\text{NH}_3)$

$^{13}\text{N}$  was generated by irradiation of 18.6 mg of  $^{13}\text{C}$  with protons (1, 32). To form  $^{13}\text{N}_2$ , the  $^{13}\text{C}$  target was subjected to Dumas combustion (1, 32). Alternatively,  $^{13}\text{N}(\text{NH}_3)$  was generated, by a modification of the method of Carangal and Varner (2, and J.E. Varner, personal communication). The target was transferred to a 100-ml Kjeldahl flask preheated to 100°C on a heating mantle. Potassium dichromate, 0.4 g, and 0.2 g potassium sulfate were added to the flask and mixed well with the target, and 2 ml of concentrated sulfuric acid (specific gravity 1.84) saturated with potassium dichromate were then added. The temperature in the flask rose steadily, reaching nearly 250°C at the end of 10 min. The flask was then removed from the heating mantle, and 5 ml of saturated sodium borate (pH 10), 8 ml of 40 percent (w/v) NaOH and 2 ml of saturated silver sulfate solution were added in rapid succession.  $^{13}\text{N}(\text{NH}_3)$  was distilled under vacuum, at 50°C, from the resulting blackish green alkaline reaction mixture. A side-arm test-tube immersed in liquid nitrogen served as receiving tube. Distillation of  $^{13}\text{N}(\text{NH}_3)$ , monitored with a model ABC-10KG-SB ionization gauge (Jordan Electronics, Division of Victoreen Instruments Co., Alhambra, Ca.), is complete within 10 to 15 s after application of vacuum from a tap-water aspirator. Bumping and

broken, but a large number of short filaments still remained intact. Twenty to twenty-five ml of the suspension were then subjected to sonic disruption under Ar for periods of 6 s.ml<sup>-1</sup> (on occasion up to 14 s.ml<sup>-1</sup>) at setting 4 of a Sonifier Cell Disruptor (Model W 185, Heat Systems-Ultrasonic Inc., Plainview, N.Y.). The suspension was then centrifuged for 10 min at 200 x g in degassed, 15-ml conical centrifuge tubes. The pellets were washed twice with, and finally resuspended in 2 to 5 ml of, a solution containing 10 mM sodium dithionite and 2 or 5 mM TES. For certain experiments, dithionite was either omitted from all solutions, or was replaced by sodium ascorbate (final concentration, 2.5 or 5 mM) throughout.

The number of heterocysts per ml of suspensions of whole filaments and of isolated heterocysts was determined with an eosinophil counter. Suspensions of isolated heterocysts (Fig. 1) contained apparently intact heterocysts as well as heterocysts which were apparently damaged (empty-looking, or with broken walls or displaced polar bodies). The sum of such "intact" and "damaged" heterocysts is referred to as "total" heterocysts. No more than two percent as many vegetative cells as heterocysts were present together with the heterocysts isolated by the procedure described above.

#### Acetylene reduction assay

Acetylene reduction by whole filaments of Anabaena was determined as described in (35), under a gas phase of Ar/C<sub>2</sub>H<sub>2</sub> (9:1, v/v), after storage of the concentrated suspension under Ar for 3 to 4.5 h, during which time the process of isolating heterocysts was completed. Comparisons are thus made between suspensions of filaments and heterocysts with the same history of exposure to aerobic and anaerobic conditions.

"boiling over" of the reaction mixture is prevented by permitting a feeble stream of air to enter the distillation apparatus (28). The frozen solution of  $^{13}\text{N}]\text{NH}_3$ , containing an average of 6 mCi  $^{13}\text{NH}_3$  and approximately 0.3  $\mu\text{moles NH}_3$ , was thawed and diluted to 1-2 ml. Samples of 25 to 100  $\mu\text{l}$  were taken for measurement of radioactivity by liquid scintillation spectrometry (26, 34).

Fixation of  $^{13}\text{N}]\text{N}_2$  was performed in the presence of Ar, 1%  $\text{CO}_2$  (26, 34), or following addition of 0.1 atm acetylene. The supplements specified earlier for assays of acetylene reduction, and sometimes 1 mM glutamate, were added prior to addition of isolated heterocysts to the reaction vial. Suspensions, 0.25 ml, of filaments (27  $\mu\text{g}$  Chl $\cdot\text{ml}^{-1}$ , without supplements) or heterocysts were exposed to  $^{13}\text{N}]\text{N}_2$  for 0.25 min or 2 (sometimes 15) min, respectively, in the presence of 4000 lux of light from an incandescent bulb. At the end of exposure, the suspensions were mixed with four volumes of methanol (26, 34). Ammonia and glutamine amide nitrogen in the methanolic extracts were separately distilled and their content of  $^{13}\text{N}$  determined with a scintillation counter (26, 34). Alternatively, the radioactive organics in the methanolic extracts were separated by high voltage (3000 v) electrophoresis with 70 mM borate buffer, pH 9.2, on thin layers of cellulose (26, 34). Radioactive constituents were localized and quantified by radiochromatogram scanning, and identified, as described previously (26, 34).

Experiments using  $^{13}\text{N}]\text{NH}_3$  were done with heterocysts isolated

in the presence of either dithionite or ascorbate, and incubated as described in Table 6. When the rate of formation of glutamine was to be quantified as nmoles  $\cdot (10^6 \text{ heterocysts})^{-1} \cdot \text{min}^{-1}$ , 2 mM  $\text{NH}_4\text{Cl}$  was also added. When evidence was sought for the occurrence of glutamate synthase activity in non-broken heterocysts, the reaction mixture in the vials was also supplemented with  $\alpha$ -ketoglutarate (up to 5 mM), and sometimes glutamine (up to 10 mM), or combinations of these. Electron donors tried were either 0.5 mM methyl viologen plus 10 mM  $\text{Na}_2\text{S}_2\text{O}_4$ , or 50  $\mu\text{M}$  DCFIP (2,6-dichlorophenol indophenol) plus 5 mM ascorbate, in the light. After 10 min of illumination, the suspensions were processed for electrophoresis (26, 34), with heterocyst lipids first displaced from  $^{13}\text{N}$ -labeled constituents by thin layer chromatography in chloroform/methanol, 1.3:1 (v/v).

The suspension of heterocysts was sometimes sedimented at 1000 x g before extraction with methanol, and the supernatant and pellet fractions processed separately for electrophoretic examination of labeled products. Where indicated, the radioactive compound co-electrophoresing with stable glutamine was eluted from the thin layer plate. The  $^{13}\text{N}$ -radioactivity of its amide nitrogen was then determined by steam distillation in the presence of alkali, and scintillation counting of the distillate (26, 34).

Labeling with  $^{14}\text{C}]\text{glutamate}$  and  $^{14}\text{C}]\alpha\text{-ketoglutarate}$

Formation of  $^{14}\text{C}$ -labeled glutamine from  $^{14}\text{C}$ -labeled glutamate (New England Nuclear; 230  $\mu\text{Ci}/\mu\text{mole}$ ) by isolated heterocysts was

determined in the presence of ATP, an ATP generating system, and  $N_2$ ,  $NH_3$ , or no exogenous nitrogen source (details in Table 5). Formation of  $^{14}C$ -labeled glutamate from  $^{14}C$ -labeled  $\alpha$ -ketoglutarate (New England Nuclear; 255  $\mu Ci/\mu mole$ ) by heterocysts in the absence of any added electron carrier was determined with various amino acids as nitrogen donor, and with or without 5 mM aminoxy acetate (an inhibitor of aminotransferases: 12, 14) or 1 mM azaserine (an inhibitor of glutamine amide transferases: 17, 34; details in Table 7).

Assays using  $^{14}C$ -labeled substrates were performed at 30°C for 30 min in 1-ml Reactivials (Pierce Chemical Co., Rockford, Ill.), with magnetic stirring. The reactions were terminated by addition of methanol and the methanolic extracts processed for electrophoresis as described above. Radioactive spots were scanned, visualized, and quantified as described earlier (26, 34).

#### Enzyme assays in vitro

Fermentor cultures, 0.6 to 1.4  $\mu g$  Chl  $\cdot ml^{-1}$ , harvested at 40,000 x g and washed with 5 mM TES, pH 7.2, were resuspended at 90 to 125  $\mu g$  Chl  $\cdot ml^{-1}$  with the same buffer. The suspension was degassed, and

supplemented with  $Na_2S_2O_4$  and dithiothreitol to final concentrations of 2.5 and 1.0 mM, respectively. For breakage of all vegetative cells and essentially all heterocysts, the algal suspension was cavitated at 12°C under argon at 3.0 amperes output of a Model S-125 sonifier (Heat Systems) for 60  $s \cdot ml^{-1}$  of suspension (29). The cavitated preparations were clarified by centrifugation at 5,000 x g for 5 min.

For studies on the time course of solubilization of glutamine synthetase and glutamate synthase, 12-ml samples which had been cavitated for the desired number of  $s \cdot ml^{-1}$  were centrifuged anaerobically at 5000 x g for 5 min. The supernatant fluid was assayed for enzymic activities (see below). The pellets, resuspended with 12 ml of buffer containing 5 mM TES, 2.5 mM  $Na_2S_2O_4$ , and 1.0 mM dithiothreitol, were cavitated for an additional 60  $s \cdot ml^{-1}$ . These suspensions were again clarified by centrifugation, and their supernatant fluid assayed.

Glutamine synthetase activity was determined by the formation of  $^{14}C$ -labeled glutamine from  $^{14}C$ -labeled glutamate as with isolated heterocysts. The reaction mixture contained, in a final volume of 0.2 ml: 5 mM ATP, 15 mM  $MgCl_2$ , 5 mM  $NH_4Cl$ , 2.5 mM glutamate (0.92 to 1.66  $\mu Ci$   $^{14}C$ ), 20 mM TES, pH 7.2, and extract containing 0.06-0.33 mg protein. Activity in the crude extract was proportional to time for 30 min, and proportional to protein concentration to 1.75 mg protein  $\cdot ml^{-1}$ .

Glutamate synthase activity was determined by the formation of  $^{14}C$ -labeled glutamate from  $^{14}C$ -labeled  $\alpha$ -ketoglutarate and stable glutamine in the presence of aminoxy acetate (11). The reaction mixture contained in a final volume of 0.2 ml: 5 mM glutamine, 1 mM methyl viologen, 12.5 mM  $Na_2S_2O_4$ , 2.5 mM  $\alpha$ -ketoglutarate (0.92-1.73  $\mu Ci$   $^{14}C$ ), 5 mM aminoxy acetate, 28 mM TES, pH 7.2, and extract

containing 0.20-0.62 mg protein. Activity was linear with time to 30 min; and was a linear function of protein concentration in the range 0.15 mg protein (essentially nil activity) to 1.0 mg protein per assay.

#### RESULTS

##### Acetylene reduction by isolated heterocysts

Heterocysts isolated in the presence of TES, sodium dithionite, EDTA and mannitol accounted normally for about 13 percent, but up to 34 percent, of the acetylene reducing activity of whole filaments (Table 1: Expt. 1). The maximum percent activity measured was 41 if only "intact" heterocysts were considered. The activity of  $11.9 \text{ nmoles } C_2H_4 \cdot (10^6 \text{ total heterocysts})^{-1} \cdot (20 \text{ min})^{-1}$  (Table 1: Expt. 1) corresponds, on the basis of 0.30 pg Chl per heterocyst (33), to an activity of  $1560 \text{ nmoles } C_2H_4 \cdot \text{mg Chl}^{-1} \cdot \text{min}^{-1}$ . Incubation in the presence of TES-dithionite-EDTA-mannitol for 45 min yielded active heterocysts. After only 5 min of incubation in this mixture, sonic treatment

resulted in preparations of heterocysts with very low nitrogenase activity (Table 1: Expt. 2), perhaps due to lesser permeation of cofactors.

Optimum acetylene reducing activity of isolated heterocysts was obtained in the presence of ATP, an ATP-generating system, and dithionite (Table 2; cf. ref. 21). Although ATP itself was required for reduction of acetylene (Table 2), acetylene was reduced at a constant but lower rate in the absence of an ATP generating system

(Figure 2, Table 2).

Replacement of sodium dithionite (13 experiments) with sodium ascorbate (9 experiments) during the isolation procedure resulted in heterocysts with, on the average, lower acetylene-reducing activity (Table 2). However, we have not found a physiological substance which could replace dithionite as an electron donor during the acetylene reduction assay. Heterocysts isolated in the absence of a reductant, and assayed in the presence of dithionite, had less than 4 percent of the activity of heterocysts isolated in the presence of dithionite. Ascorbate plus DCPIP could not replace dithionite as an electron donor in the light (Table 2).

##### Fixation of $^{13}N$ -labeled nitrogen gas and coupling to glutamine formation, by isolated heterocysts

Heterocysts isolated in the presence of dithionite or ascorbate reduced  $[^{13}N]N_2$  in the light. Scans of electrophoretograms of methanolic extracts showed only a single organic product of fixation of  $[^{13}N]N_2$ , which co-electrophoresed with  $^{14}C$ -labeled glutamine (Fig. 3). Vacuum distillation at pH 10 of methanolic extracts of heterocysts labeled for 120 s with  $[^{13}N]N_2$  was followed by steam-distillation in the presence of alkali. These distillation procedures showed the presence of  $[^{13}N]NH_3$  and  $^{13}N$ -labeled amide nitrogen (Table 3; see 26, 34). Addition of 0.1 atm of acetylene to the reaction vial containing heterocysts inhibited  $[^{13}N]N_2$  fixation by about 99 percent.

Radioactivity of the amide nitrogen steam-distilled as described above accounted for up to 67 percent of the combined distillates for ammonia and amide nitrogen after fixation of  $[^{13}N]N_2$  for 120 s by heterocysts in the presence of 1 mM glutamate (Table 3). Radioactivity

in the combined distillates of extracts of heterocysts accounted for up to 23 percent of the label found in the combined distillates ( $\text{NH}_3$  and amide-N) of extracts of whole filaments exposed to [ $^{13}\text{N}$ ] $\text{N}_2$ , per heterocyst and per min of fixation (Table 4), or for about 20 percent of the  $^{13}\text{N}$ -radioactivity present in whole filaments, if  $\alpha$ -amino- $^{13}\text{N}$  of glutamate and glutamine is considered (see ref. 34). We used only 15-s exposures of filaments to [ $^{13}\text{N}$ ] $\text{N}_2$  so that the great majority of fixed  $^{13}\text{N}$  would be distillable, and longer exposure of heterocysts so as to increase fixation, and thereby reduce the statistical error in determination of the radioactivity fixed. Examination, under a microscope, indicated that approximately 50 percent of the population of isolated heterocysts used in this experiment could be considered "damaged." If  $\text{N}_2$ -fixing activity were restricted to "intact" heterocysts, the nitrogenase activity in such heterocysts would have accounted for about 40 percent of the  $\text{N}_2$ -fixing activity of whole filaments.

When isolated heterocysts were incubated in the light for 30 min in the presence of  $^{14}\text{C}$ -labeled glutamate and other supplements, and with either  $\text{N}_2$  or no other added source of nitrogen, about 22 percent as much [ $^{14}\text{C}$ ]glutamine was formed as in the presence of 5 mM ammonia (Table 5). Addition of 1 mM methionine sulfoximine inhibited formation of  $^{14}\text{C}$ -labeled glutamine by nearly 90 percent.

#### Assimilation of $^{13}\text{N}$ -labeled ammonia by isolated heterocysts

Isolated heterocysts formed, from [ $^{13}\text{N}$ ] $\text{NH}_3$ , a  $^{13}\text{N}$ -labeled substance which co-electrophoresed with stable and with  $^{14}\text{C}$ -labeled glutamine. Addition of stable glutamate to the reaction mixture resulted in greatly increased radioactivity of  $^{13}\text{N}$ -labeled glutamine (Table 6). When the region of the thin layer plate bearing  $^{13}\text{N}$ -radioactivity was

eluted, and the eluate steam-distilled in the presence of alkali (26, 34), all of the original radioactivity, within experimental error, could be accounted for as amide- $^{13}\text{N}$ . The average rate of formation of glutamine observed in the presence of 1 mM glutamate and 2 mM  $\text{NH}_4\text{Cl}$  was 0.1  $\mu\text{moles} \cdot (10^6 \text{ heterocysts})^{-1} \cdot \text{min}^{-1}$ , whereas  $\text{N}_2$ -fixing filaments supplemented with 1 mM glutamate and 2 mM  $\text{NH}_4\text{Cl}$  formed an average of 10  $\mu\text{moles} \text{ glutamine} \cdot (10^6 \text{ heterocysts})^{-1} \cdot \text{min}^{-1}$ . When the suspension of heterocysts in the reaction vial was sedimented, after incubation with [ $^{13}\text{N}$ ] $\text{NH}_3$ , glutamate, and other cofactors, and the pellet and supernatant fluid processed separately, 97 percent of the radioactive glutamine was found in the supernatant fluid (Table 6). Thoroughly extracted pellets also contained slight radioactivity. In comparable experiments in which isolated heterocysts were incubated with [ $^{14}\text{C}$ ]glutamate and  $\text{NH}_3$ , the [ $^{14}\text{C}$ ]glutamine which was formed and the [ $^{14}\text{C}$ ]glutamate which remained were both distributed 0.96:0.04 between the supernatant and pellet fractions. In the presence of 1 mM methionine sulfoximine, very little (Table 6)  $^{13}\text{N}$ -labeled substance detectable by scanning of electrophoretograms was formed.

#### Localization of glutamate formation

Heterocysts were incubated for 10 min with [ $^{13}\text{N}$ ] $\text{NH}_3$  and other cofactors (see above) for synthesis of [ $^{13}\text{N}$ ]glutamine. Adding  $\alpha$ -ketoglutarate (up to 5 mM), glutamine (up to 10 mM), and methyl viologen plus dithionite or DCPIP plus ascorbate in the light as electron donor, and decreasing the concentration of glutamate from 1 mM to 0.1 or 0 mM, resulted in a ratio of [ $^{13}\text{N}$ ]glutamate to [ $^{13}\text{N}$ ]glutamine of at most 0.02. These results provided no evidence for the idea that heterocysts have substantial glutamate synthase activity.



We therefore attempted to localize the site of glutamate synthase activity by the technique of differential disruption of vegetative cells and heterocysts (29, 36). Enzymatic activities associated with supernatant and pellet fractions after cavitation for periods of 1 to 25 s·ml<sup>-1</sup> of a suspension of filaments were assayed in cell-free extracts as described in Materials and Methods. As previously shown (29), the number of apparently intact heterocysts decreases exponentially during cavitation, with 50 percent of the heterocysts remaining apparently intact following cavitation for about 15 s·ml<sup>-1</sup> of suspension, whereas the amounts of chlorophyll (Fig. 4) and protein (not shown) sedimentable at 5000 x g show a much more rapid decrease. Glutamate synthase activity is solubilized very rapidly from the filaments, in that fifty percent of the glutamate synthase activity is found in the supernatant fluid following cavitation for close to 1 s·ml<sup>-1</sup> of suspension, and essentially 100 percent after cavitation for 5 s·ml<sup>-1</sup> of suspension. Vegetative cells are destroyed comparably rapidly (36). After 5 s·ml<sup>-1</sup> of cavitation, and centrifugation for 5 min at 5000 x g, 75 to 80 percent of the heterocysts remained apparently intact and were sedimented, whereas approximately 25 percent of the glutamate synthetase activity and approximately 22 percent of the total protein and chlorophyll were sedimented. Cavitation for up to 25 s·ml<sup>-1</sup> of suspension continued to solubilize glutamate synthetase activity, approximately in parallel with the "solubilization" of chlorophyll (Fig. 4) and protein (not shown).

#### Aminotransferase activity of isolated heterocysts

Isolated heterocysts, incubated for 30 min in the presence of and <sup>14</sup>C-labeled α-ketoglutarate, formed [<sup>14</sup>C]glutamate at

a substantial rate (Table 7). This activity was extensively inhibited by 5 mM aminoxy acetate, but was not affected by 1 mM azaserine. The formation of [<sup>14</sup>C]glutamate, also unlike the glutamate synthase reaction in vitro, was not dependent on the presence of an exogenous reductant such as dithionite-reduced methyl viologen. The aminotransferase activity could also be supported by glutamine, at a lower rate than by glutamate, and by aspartate, but not measurably by alanine. Glutamate-supported transferase activity was also unaffected by azaserine and inhibited by aminoxy acetate. Some formation of [<sup>14</sup>C]glutamate from <sup>14</sup>C-labeled α-ketoglutarate occurred in the absence of exogenous amino donors.

#### DISCUSSION

##### Nitrogenase activity of heterocysts

Our results with isolated heterocysts provide additional direct evidence that these cells are major sites of nitrogen fixation during aerobic photosynthetic growth of filamentous cyanobacteria. Fixation of N<sub>2</sub> by isolated heterocysts had not heretofore been demonstrated. The isolated heterocysts in our experiments accounted for a higher fraction of intact-filament nitrogenase activity than did the heterocysts isolated from a different organism by Peterson and Burris (16), but were of slightly (ca. 15%) lower specific activity, and had a far higher specific activity than that reported (23) for heterocysts isolated from a different strain of Anabaena cylindrica. Hydrogen, included in the gas phase over the suspensions and in the assay vials used by Peterson and Burris (16), may have stimulated acetylene reduction activity (cf. 35). Our procedure, designed to obtain heterocyst preparations that would reduce [<sup>13</sup>N]N<sub>2</sub> as well as acetylene, omitted hydrogen because of its

inhibitory effect on  $N_2$  fixation.

"Intact" heterocysts in our suspensions accounted for up to 40 percent of the nitrogenase activity of intact filaments. The initial localization of the remaining activity is unknown. The results of autoradiographic experiments with [ $^{13}N$ ]  $N_2$  (32) and of experiments on the time course of solubilization of nitrogenase during cavitation (36) were consistent with, but did not prove, the interpretation that the remaining activity is located in the vegetative cells. However, sonic disruption is known to damage heterocysts (5). By preincubating in TES-dithionite-EDTA-mannitol we have been able to reduce the time of sonic disruption required to isolate heterocysts to  $6 \text{ s} \cdot \text{ml}^{-1}$ . This much cavitation may, however, still damage the functionality of the heterocysts. In the absence of a reliable procedure for evaluating the proportion of truly intact, isolated heterocysts, it is difficult to arrive at a firm conclusion about the percent of nitrogenase located in heterocysts, in vivo.

#### Enzymes involved in assimilation of $NH_3$ by heterocysts

Dharmawardene et al. (3) showed that the specific activity of glutamine synthetase in isolated heterocysts is 1.7 times greater than in vegetative cells. We have found that after  $12.5$  or  $25 \text{ s} \cdot \text{ml}^{-1}$  of sonic disruption, the ratio of sedimentable glutamine synthetase activity per intact heterocyst (no vegetative cells remain intact), corrected to the initial number of heterocysts, accounts for about 8% of the initial activity of glutamine synthetase (Fig. 4). This figure is slightly in excess of the initial ratio of heterocysts to total cells. We have thus confirmed that in contrast to enzymes of the oxidative pentose phosphate cycle (13, 19, 29), there is not a great difference in the specific activity of glutamine synthetase in heterocysts compared with vegetative cells. A number of observations, taken together, show that glutamine synthetase is involved in the assimilation of ammonia by heterocysts. First, radioactive glutamine is formed from [ $^{13}N$ ]  $NH_3$  and from [ $^{14}C$ ] glutamate, the synthesis being in

both cases largely inhibited by the addition of methionine sulfoximine. We have not attempted to account for the limited formation of [ $^{14}C$ ] glutamine in the presence of methionine sulfoximine. Second, synthesis of [ $^{13}N$ ] glutamine is enhanced by addition of glutamate, and synthesis of [ $^{14}C$ ] glutamine is enhanced by addition of ammonia. Finally, the [ $^{13}N$ ] in the glutamine synthesized was all, or nearly all, in the amide group.

Moreover, using [ $^{13}N$ ]  $N_2$  we have demonstrated that nitrogenase is indeed coupled metabolically to glutamine synthetase in heterocysts. Addition of 0.1 atm of acetylene, an inhibitor of nitrogen fixation (18), decreased the [ $^{13}N$ ] radioactivity in ammonia plus glutamine by about 99 percent, indicating that glutamine is formed from  $N_2$ -derived ammonia. Because formation of [ $^{14}C$ ] glutamine from [ $^{14}C$ ] glutamate is little affected by substitution of Ar for  $N_2$  in the gas phase (Table 5), an endogenous source of  $NH_3$  appears also to be present in the suspensions. Although isolated heterocysts were observed to metabolize two-thirds of their newly fixed nitrogen to glutamine, this figure may underestimate the in-vivo fraction. That is, if the isolation procedure increases the "leakiness" of heterocysts to substances of low molecular weight,  $N_2$ -derived [ $^{13}N$ ]  $NH_3$  may leak out of isolated heterocysts before it can react with glutamine synthetase.

When heterocysts were incubated with [ $^{13}N$ ]  $NH_3$  and stable or [ $^{14}C$ ]  $NH_3$ -labeled glutamate, the suspension sedimented, and the supernatant fluid and pellet examined separately for (extractable) radioactivity, it was found that 96-97 percent of the radioactive glutamine formed was in the supernatant fluid. Thus, the glutamine synthesized in the isolated heterocysts is rapidly lost from those cells.

In extracts of  $N_2$ -grown filaments (11), but not in suspensions of heterocysts,  $\alpha$ -ketoglutarate is aminated to form glutamate in the presence of glutamine, dithionite and methyl viologen.

Possible implications for pattern formation

It is known that heterocysts inhibit nearby cells in the same filament from differentiating into heterocysts (30, 31), that the inhibition is mediated by a diffusible inhibitor produced by heterocysts (31), and that methionine sulfoximine, which inhibits synthesis of glutamine from ammonia (22, 34) prevents exogenously supplied ammonia (9) from inhibiting heterocyst formation (22). It is therefore possible that the inhibitor which moves from cell to cell may be glutamine or a derivative of glutamine. If this suggestion were correct, it might be that either glutamine itself, or some derivative of it, is directly involved, in vegetative cells, in inhibiting differentiation.

ACKNOWLEDGMENTS

This research was supported by the U.S. Energy Research and Development Administration under Contract E(11-1)-1338, and by the National Science Foundation.

Massive accumulation of  $^{13}\text{N}$ -radioactivity in glutamine was observed in suspensions of heterocysts (Table 6) with very little or no appearance of radioactivity in glutamate. [ $^{14}\text{C}$ ]labeled  $\alpha$ -ketoglutarate easily enters isolated heterocysts, as shown by its extensive transamination to form glutamate. This reaction is indeed a transamination, because it is greatly inhibited by aminoxy acetate (12, 14). Dithionite and ATP are presumed to enter heterocysts, because of the dependence of nitrogenase activity on their presence. However, whether methyl viologen, ascorbate and DCPIP can penetrate the heterocysts is unknown.

We therefore tested by means of in-vitro reactions whether glutamate synthase activity is concentrated in heterocysts. Our results (Fig. 4) appear to show that cavitation renders glutamate synthase activity non-sedimentable much more rapidly than it does glutamine synthetase activity, which is in turn much more rapidly solubilized than are enzymes of the oxidative pentose phosphate pathway (29). Accordingly, we suggest that the specific activity of glutamate synthase, the third enzyme in the initial pathway of assimilation of  $\text{N}_2$  in *Anabaena cylindrica* (26, 34), is much greater in vegetative cells than in heterocysts. It therefore appears that glutamine newly synthesized in heterocysts is transported to vegetative cells, where glutamate is synthesized. It further appears that part of the glutamate formed then returns to the heterocysts to be amidated or, alternatively, that  $\alpha$ -ketoglutarate and amidated substrates of the aminotransferase of heterocysts react within heterocysts to form glutamate which is then amidated.

9. Fogg, G.E. 1949. Growth and heterocyst production in Anabaena cylindrica Lemm. II. In relation to carbon and nitrogen metabolism. Ann. Bot. N.S. 13:241-259.
10. Haystead, A.R., R. Robinson, and W.D.P. Stewart. 1970. Nitrogenase activity in extracts of heterocystous and non-heterocystous blue-green algae. Arch. Mikrobiol. 74:235-243.
11. Lea, P.J., and B.J. Mifflin. 1975. Glutamate synthase in blue-green algae. Biochem. Soc. Trans. (U.K.) 3:381-384.
12. Lea, P.J., and B.J. Mifflin. 1975. The occurrence of glutamate synthase in algae. Biochem. Biophys. Res. Commun. 64:856-862.
13. Lex, M., and N.G. Carr. 1974. The metabolism of glucose by heterocysts and vegetative cells of Anabaena cylindrica. Arch. Mikrobiol. 101:161-167.
14. Mifflin, B.H., and P.J. Lea. 1975. Glutamine and asparagine as nitrogen donors for reductant-dependent glutamate synthesis in pea roots. Biochem. J. 149:403-409.
15. Nagatani, H., M. Shimizu, and R.C. Valentine. 1971. The mechanism of ammonia assimilation in nitrogen-fixing bacteria. Arch. Mikrobiol. 79:164-175.
16. Peterson, R.B., and R.H. Burris. 1976. Properties of heterocysts isolated with colloidal silica. Arch. Mikrobiol. 108:35-40.
17. Prusiner, S., and E.R. Stadtman. 1973. The enzymes of glutamine metabolism, Academic Press, New York.
18. Rivera-Ortiz, J.M., and R.H. Burris. 1975. Interactions among substrates and inhibitors of nitrogenase. J. Bacteriol. 123:537-545.
19. Rowell, P., and W.D.P. Stewart. 1976. Alanine dehydrogenase of the N<sub>2</sub>-fixing blue-green alga, Anabaena cylindrica. Arch. Mikrobiol. 107:115-124.

## LITERATURE CITED

1. Austin, S.M., A. Galonsky, J. Bortins, and C.P. Wolk. 1975. A batch process for the production of <sup>13</sup>N-labeled nitrogen gas. Nuclear Inst. & Meth. 126:373-379.
2. Carangal, A.R., and J.E. Varner. 1959. The preparation of <sup>13</sup>N-labeled compounds for nitrogen metabolism studies. Plant Physiol. (Suppl.) 34:xi.
3. Dharmawardene, M.W.N., A. Haystead, and W.D.P. Stewart. 1973. Glutamine synthetase of the nitrogen-fixing alga Anabaena cylindrica. Arch. Mikrobiol. 90:281-295.
4. Fay, P. 1969. Cell differentiation and pigment composition in Anabaena cylindrica. Arch. Mikrobiol. 67:62-70.
5. Fay, P., and N.J. Lang. 1971. The heterocysts of blue-green algae. I. Ultrastructural integrity after isolation. Proc. Roy. Soc. B 178:185-192.
6. Fay, P., W.D.P. Stewart, A.E. Walsby, and G.E. Fogg. 1968. Is the heterocyst the site of nitrogen fixation in blue-green algae? Nature (London) 220:810-812.
7. Fay, P., and A.E. Walsby. 1966. Metabolic activities of isolated heterocysts of the blue-green alga Anabaena cylindrica. Nature (London) 209:94-95.
8. Fleming, H., and R. Haselkorn. 1973. Differentiation in Nostoc muscorum: Nitrogenase is synthesized in heterocysts. Proc. Nat. Acad. Sci. U.S.A. 70:2727-2731.

20. Smith, R.V., and M.C.W. Evans. 1970. Soluble nitrogenase from vegetative cells of the blue-green alga Anabaena cylindrica. Nature (London) 225:1253-1254.
21. Stewart, W.D.P., A. Haystead, and H.W. Pearson. 1969. Nitrogenase activity in heterocysts of filamentous blue-green algae. Nature (London) 224:226-228.
22. Stewart, W.D.P., and P. Rowell. 1975. Effects of L-methionine-DL-sulphoximine on the assimilation of newly fixed  $\text{NH}_3$ , acetylene reduction and heterocyst production in Anabaena cylindrica. Biochem. Biophys. Res. Commun. 65:846-856.
23. Tel-Or, E., and W.D.P. Stewart. 1976. Photosynthetic electron transport, ATP synthesis and nitrogenase activity in isolated heterocysts of Anabaena cylindrica. Biochim. Biophys. Acta 423:189-195.
24. Tempest, D.W., J.L. Meers, and C.M. Brown. 1970. Synthesis of glutamate in Aerobacter aerogenes by a hitherto unknown route. Biochem. J. 117:405-407.
25. Thomas, J., and K.A.V. David. 1972. Site of nitrogenase activity in the blue-green alga Anabaena sp. L-31. Nature New Biol. (London) 238:219-221.
26. Thomas, J., C.P. Wolk, P.W. Shaffer, S.M. Austin, and A. Galonsky. 1975. The initial organic products of fixation of  $^{13}\text{N}$ -labeled nitrogen gas by the blue-green alga Anabaena cylindrica. Biochem. Biophys. Res. Commun. 67:501-507.
27. Van Gorkom, H.J., and M. Donze. 1971. Localization of nitrogen fixation in Anabaena. Nature (London) 234:231-232.

28. Varner, J.E., W.A. Eulen, S. Vanecko, and R.C. Burrell. 1953. Determination of ammonium, amide, nitrite and nitrate nitrogen in plant extracts. Anal. Chem. 25:1528-1529.
29. Winkenschach, F., and C.P. Wolk. 1973. Activities of enzymes of the oxidative and/reductive pentose phosphate pathways in heterocysts of a blue-green alga. Plant Physiol. 52:480-483.
30. Wolk, C.P. 1967. Physiological basis of the pattern of vegetative growth of a blue-green alga. Proc. Nat. Acad. Sci. U.S.A. 57:1246-1251.
31. Wolk, C.P. 1975. Formation of one-dimensional patterns by stochastic processes and by filamentous blue-green algae. Dev. Biol. 46:370-382.
32. Wolk, C.P., S.M. Austin, J. Bortins, and A. Galonsky. 1974. Autoradiographic localization of  $^{13}\text{N}$  after fixation of  $^{13}\text{N}$ -labeled nitrogen gas by a heterocyst-forming blue-green alga. J. Cell Biol. 61:440-453.
33. Wolk, C.P., and R.D. Simon. 1969. Pigments and lipids of heterocysts. Planta (Berlin) 86:92-97.
34. Wolk, C.P., J. Thomas, P.W. Shaffer, S.M. Austin, and A. Galonsky. 1976. The pathway of nitrogen metabolism after fixation of  $^{13}\text{N}$ -labeled nitrogen gas by the cyanobacterium Anabaena cylindrica. J. Biol. Chem. In press.
35. Wolk, C.P., and E. Wojciuch. 1971. Photoreduction of acetylene by heterocysts. Planta (Berlin) 97:126-134.
36. Wolk, C.P., and E. Wojciuch. 1971. Biphasic time course of solubilization of nitrogenase during cavitation of aerobically grown Anabaena cylindrica. J. Phycol. 7:339-344.

Table 1. Acetylene reduction by isolated heterocysts and whole filaments of *Anabaena cylindrica*

Expt. No.	Conditions for isolation of heterocysts	nmoles $C_2H_4$ formed per 20 min <sup>a</sup>	
		per 10 <sup>6</sup> isolated heterocysts	per 10 <sup>6</sup> "total" heterocysts of whole filaments <sup>c</sup>
1.	Normal procedure: incubation for 45 min in 1.4 mM TES, 6.5 mM $Na_2S_2O_4$ , 26 mM $Na_2EDTA$ , 650 mM D-mannitol	11.9 <sup>d</sup> (34)	14.3 (41)
2.	As in 1, but only 5 min incubation	0.7 (1.5)	47.7(100)

a. Acetylene reduced at 30°C under 4000 lux incandescent light in 5-ml serum vials containing 1.8-1.9 x 10<sup>6</sup> heterocysts isolated in the presence of dithionite, or containing suspensions of whole filaments having 0.94-2.3 x 10<sup>6</sup> heterocysts. Vials containing isolated heterocysts were supplemented with 0.5 mg creatine phosphokinase, 50 mM creatine phosphate, 5 mM ATP, 5 mM MgCl<sub>2</sub>, 10 mM  $Na_2S_2O_4$  and 50 mM TES in 2 ml final volume. Gas phase: 90% Ar, 10%  $C_2H_2$ . All values are means of duplicate determinations. Values in parentheses are percentages relative to intact filaments.

b. "Intact" and "total" heterocysts are defined in Materials and Methods.

c. Filaments concentrated to 27-100  $\mu g$   $Chl \cdot ml^{-1}$  were stored under Ar for 3 to 4.5 h until the isolation of heterocysts was completed, and were then assayed for reduction of  $C_2H_2$ .

d. Not determined.

Table 2. Requirements for acetylene reduction in the light by heterocysts isolated in the presence or absence of sodium dithionite or sodium ascorbate

Reductant present during isolation: Assay conditions	nmoles $C_2H_4$ formed per 10 <sup>6</sup> total isolated heterocysts in 20 min	
	10 mM Dithionite	5 mM Ascorbate
Complete supplements <sup>a</sup>	13.3	3.74
$-Na_2S_2O_4$ + 5 mM sodium ascorbate + 50 $\mu M$ DCPiP	0.40	0.06
-Creatine phosphate (CrP)	9.4	2.35
-creatine phosphokinase (CrK)	0.0	0.49
-CrP-CrK-ATP	0.0	0.23

a. As in Table 1.

b. Not determined.

Table 3. Distribution of  $^{13}\text{N}$ -radioactivity in products distilling as ammonia and amide nitrogen after fixation of  $^{13}\text{N}_2$  by heterocysts isolated from *Anabaena cylindrica*

	cpm $\mu\text{Ci}^{-1}$ <sup>a</sup>	
	$^{13}\text{N}:\text{NH}_3$	Amide- $^{13}\text{N}$
Standard <sup>b</sup> , + heterocysts	15.93±0.29 <sup>c</sup>	31.74±0.23 <sup>c</sup>
Standard, + heterocysts, + 0.1 atm $\text{C}_2\text{H}_2$	0.18±0.02	0.28±0.03
Standard, - heterocysts	-0.04±0.04	0.04±0.13

a. After fixation of  $^{13}\text{N}_2$  for 120 s, the suspension was extracted with 80% methanol and the extract subjected first to vacuum distillation at pH 10, to determine " $^{13}\text{N}:\text{NH}_3$ ", and then to steam distillation in the presence of NaOH, to determine "amide- $^{13}\text{N}$ ". Cpm  $\mu\text{Ci}^{-1}$  represents cpm  $^{13}\text{N}:\text{NH}_3$  or cpm amide- $^{13}\text{N}$  in the fixation vial (measured with approximately 74% counting efficiency), corrected to the time of the start of fixation, and normalized to equal amounts of  $^{13}\text{N}$  (in  $\mu\text{Ci}$ ) in the vial.

b. One-ml reaction vials contained, in a volume of 0.25 ml, 1 mM glutamate, 4 mM ATP, 4 mM  $\text{MgCl}_2$ , 40 mM phosphocreatine, 0.05 mg creatine phosphokinase, 10 mM  $\text{Na}_2\text{S}_2\text{O}_4$ , 24 mM TES, pH 7.2, and, as indicated,  $2.55 \times 10^6$  "total" or no heterocysts, under an atmosphere of 2%  $\text{N}_2$ , 1%  $\text{CO}_2$ , balance Ar.

c.  $\bar{x} \pm \sigma_x$ , after correction for background (approximately 18 cpm) and time of decay, calculated from replicate readings of three separate samples of  $^{13}\text{N}:\text{NH}_3$  and distillate (uncorrected values 2352 to 3564 cpm) and amide- $^{13}\text{N}$  distillate (uncorrected values 2480 to 3578 cpm).

Table 4. Comparison of rates of fixation of  $^{13}\text{N}:\text{N}_2$  by filaments and by heterocysts isolated from the same culture

	Filaments	Heterocysts <sup>a</sup>
Cpm $\mu\text{Ci}^{-1}$ (see footnote a of Table 3)	2.20	8.56
Fixation time, min	0.25	2.00
Total heterocysts per vial	$1.09 \times 10^6$	$2.27 \times 10^6$
Cpm $\mu\text{Ci}^{-1} \text{min}^{-1}$ (106 heterocysts) <sup>-1</sup>	8.07 (100%)	1.88 (23%)

a. The reaction mixture was the same as in footnote b of Table 3, but with glutamate omitted.

Table 5. Formation of [<sup>14</sup>C]glutamine by isolated heterocysts, in the light.

Assay conditions <sup>a</sup>	Percent of NH <sub>3</sub> -supported activity <sup>b</sup>
Standard	23.2(4) <sup>c</sup> ± 6.9
-ATP	7.5(3) ± 2.7
-N <sub>2</sub> + Ar	20.9(4) ± 6.9
+NH <sub>3</sub>	100.0
+NH <sub>3</sub> + MSX	10.0(3) ± 5.6

a. Reaction mixtures contained, in a volume of 0.2 ml, 5 mM ATP, 15 mM MgCl<sub>2</sub>, 50 mM creatine phosphate, 0.05 mg creatine phosphokinase, 15 mM Na<sub>2</sub>S<sub>2</sub>O<sub>4</sub>, 60 mM TES, pH 7.2, 20 to 36 μM [<sup>14</sup>C]glutamate, and 0.7 to 2.2 x 10<sup>6</sup> heterocysts under N<sub>2</sub> ("standard"). Where indicated, ATP was not included, N<sub>2</sub> was replaced with Ar, or 1mM methionine sulfoximine (MSX) and/or 5 mM NH<sub>4</sub>Cl was added.

b. Rate of NH<sub>3</sub>-supported formation of [<sup>14</sup>C]glutamine by isolated heterocysts: 3.1 ± 0.3 nmoles·(10<sup>6</sup> heterocysts)<sup>-1</sup>·(30 min)<sup>-1</sup> (4)<sup>c</sup>.

c. Number of observations.

Table 6. Effect of added glutamate and methionine sulfoximine on the incorporation of [<sup>13</sup>N]-labeled ammonia into glutamine by heterocysts isolated from *Anabaena cylindrica*

Additions to reaction vial <sup>a</sup>	dpm glutamine- <sup>13</sup> N recovered <sup>b</sup>	dpm <sup>13</sup> NH <sub>3</sub> added <sup>b</sup>	%
1. 5 mM ATP, 8 mM MgCl <sub>2</sub> , 13 mM TES, 50 μM DCFIP, 5 mM sodium ascorbate, 5 mM α-ketoglutarate.			0.032
2. As in 1, plus 1 mM Glutamate.			1.21
3. Same as in 2; heterocysts and medium separated.	0.031 <sup>c</sup>	1.16 <sup>d</sup>	
4. As in 2, plus 1 mM methionine sulfoximine.	<0.023		

a. Reaction vials each contained 2.27 x 10<sup>6</sup> heterocysts, isolated in the presence of sodium ascorbate, and supplements, in a total volume of 0.4 ml. Reactions were initiated by the addition of [<sup>13</sup>N]-labeled ammonia, 6.32 x 10<sup>8</sup> dpm<sup>b</sup>, and were incubated for 10 min in the presence of 4000 lux incandescent light.

b. Corrected to the time of distillation of [<sup>13</sup>N]NH<sub>3</sub>.

c.d. In experiment 3, the suspension of heterocysts was sedimented by centrifugation at 1000 x g, and the pellet (c), and supernatant fluid (d) were separately extracted and processed for electrophoresis.



Table 7. Aminotransferase activity of heterocysts isolated in the presence of ascorbate

Supplements	nmoles glutamate formed $\cdot (10^6 \text{ heterocysts})^{-1} \cdot 30 \text{ min}^{-1}$
5 mM Glutamate (Glu)	1.26
5 mM Glu + 5 mM Aminoxy acetate	0.01
5 mM Glu + 1 mM Azaserine	1.22
5 mM Glutamine (Gln)	0.84
5 mM Gln + 5 mM Aminoxy acetate	0.00
5 mM Gln + 1 mM Azaserine	0.94
5 mM Asparate	1.24
5 mM Alanine	0.18
None (no exogenous N-donor)	0.20

Reaction mixtures contained, in 0.2 ml: 25 mM TES, pH 7.2, 18-34  $\mu\text{M}$  [ $^{14}\text{C}$ ]-ketoglutaric acid, plus supplements as indicated. Incubation was for 30 min at 30°C.

Fig. 1. Photomicrograph of heterocysts isolated from *Anabaena cylindrica*, and active in reduction of [ $^{13}\text{N}$ ] $\text{N}_2$  and acetylene. The heterocysts are suspended in a solution containing 2.5 mM sodium ascorbate and 5 mM TES, pH 7.2. A low frequency of broken heterocysts and less than 2 percent vegetative cells are present as contaminants.  $\times 920$ .

Fig. 2. Time course of reduction of acetylene by heterocysts isolated from *Anabaena cylindrica*. Reaction vials contained  $1.75 \times 10^6$  heterocysts in a total volume of 2 ml. Details of assay, and supplements present, were as specified for Table 1 (o-o), or creatine phosphokinase and creatine phosphate were omitted ( $\Delta$ - $\Delta$ ).

Fig. 3. Scan of radioactivity from  $^{13}\text{N}$  (---) in an electrophoretogram of compounds extracted from a suspension of heterocysts isolated from *Anabaena cylindrica* in the presence of dithionite, and allowed to fix  $[^{13}\text{N}]\text{N}_2$  for 15 min at 4000 lux incandescent light. The 1-ml reaction vial contained  $7.1 \times 10^6$  heterocysts, and supplements and gas phase as stated in footnote b of Table 3. An 80% methanolic extract of the contents of the reaction vial, supplemented with ten unlabeled amino acids and  $[^{14}\text{C}]\text{glutamine}$  as markers, was applied in a thin strip (at 0) 12 cm from the positive (left-hand) end of a cellulose thin layer plate (20 x 5 cm). Lipids were displaced from the region of the origin by chromatography in chloroform-methanol (1.3:1, v/v). The extract was then subjected to electrophoresis at 3000 V for 12 min in 70 mM sodium borate buffer, pH 9.2. The plate was scanned at  $5 \text{ cm} \cdot \text{min}^{-1}$ ,  $r=1s$ . After decay of the  $^{13}\text{N}$ , the plate was rescanned as before (---). A 127- $\mu\text{m}$  layer of aluminum covering the detector was then removed, and the electrophoretogram scanned for radioactivity from  $^{14}\text{C}$  (----). The plate was then dried at  $90^\circ\text{C}$ , and sprayed with a solution of ninhydrin to localize the stable marker amino acids. The peak of  $^{13}\text{N}$ -radioactivity in the figure co-electrophoresed with glutamine. Glutamate was present at the 6.7-cm position after electrophoresis.

**Fig. 4.** Time course of solubilization of glutamine synthetase and glutamate synthase activities during cavitation. Heterocysts apparently intact as percent of initial number of heterocysts (x); chlorophyll in 5 min, 5000 x g pellets as percent of total chlorophyll ( $\Delta$ ); activities of glutamine synthetase ( $\bullet$ ) and glutamate synthase ( $\blacksquare$ ) associated with such pellets (left-hand ordinate; see below) as percent of total activity; and activities of glutamine synthetase (o) and glutamate synthase ( $\square$ ) in the supernatant fluid (right-hand ordinate), assayed as described in Materials and Methods, as percent of total activity, after cavitation for varying periods of time. Total activities of glutamine synthetase and glutamate synthase were taken to be the activities of the supernatant fluid following cavitation of a cyanobacterial suspension for  $60 \text{ s}\cdot\text{ml}^{-1}$ , and centrifugation for 5 min at  $5000 \times g$ . These activities were  $25.4$  and  $5.2 \text{ } \mu\text{moles}\cdot(\text{mg protein})^{-1}\cdot\text{min}^{-1}$ , respectively. Enzymatic activities associated with the pellets were measured by resuspending the pellets, cavitating for an additional period of  $60 \text{ s}\cdot\text{ml}^{-1}$ , centrifuging as before, and assaying the supernatant fluid. The time course of solubilization of glucose-6-phosphate dehydrogenase (---), an activity associated largely with heterocysts, is repeated from an earlier paper (29), for the sake of comparison.





