


MICHIGAN STATE UNIVERSITY


CYCLOTRON LABORATORY

OBSERVATION OF M1 STRENGTH BY THE INELASTIC  
SCATTERING OF 200 MeV PROTONS

N. ANANTARAMAN, G. M. CRAWLEY, A. GALONSKY,  
C. DJALALI, N. MARTY, M. MORLET, A. WILLIS  
and J-C. JOURDAIN



MARCH 1981



Observation of M1 Strength by the Inelastic Scattering of 200

MeV Protons

N. Anantaraman, G.M. Crawley, and A. Galonsky  
Cyclotron Laboratory, Michigan State University,

East Lansing, MI 48824

and

C. Djalali, N. Marty, M. Morlet, A. Willis and J.-C. Jourdain  
Institut de Physique Nucleaire, 91406 Orsay Cedex, France

Abstract

A new resonance has been observed at very forward angles in the inelastic scattering of 200 MeV protons on  $^{90}\text{Zr}$ ,  $^{92}\text{Zr}$  and  $^{94}\text{Zr}$  at an excitation energy around 8.8 MeV in each target. The excitation energy, angular distribution and strength of this state suggests that it corresponds to the previously unseen giant M1 transition in the zirconium isotopes.

PACS numbers: 24.30.Cz, 25.40.Ep

The location of M1 strength in medium and heavy nuclei has been a long standing and major problem in nuclear structure physics.<sup>1</sup> While various theoretical estimates<sup>2,3</sup> have indicated that M1 states should exist below 12 MeV excitation energy in nuclei with  $A > 60$ , little or no significant M1 strength has been observed. This strength has been searched for in both hadron<sup>4,5</sup>, and electron<sup>6,7,8</sup> inelastic scattering but so far without success. There has been some M1 strength reported in  $^{208}\text{Pb}$ <sup>9</sup> and a mention<sup>10</sup> of strength observed in ( $e,e'$ ) on  $^{90}\text{Zr}$ , which has not been substantiated by later measurements.<sup>7</sup>

In contrast, there has been observed in recent charge exchange<sup>11-17</sup> experiments on more than twenty target nuclei, a broad resonance which is generally accepted as the giant Gamow-Teller (G-T) resonance. This transition, in which  $\Delta J^{\pi} = 1^+$ , is successfully accounted for in terms of the shell model<sup>18</sup>, and the same considerations would predict appreciable strength for a giant M1 state in the target. Hence, we have the paradoxical situation that the transition to the state in the daughter nucleus has been observed many times, but the parent has never been seen. The elusiveness of the parent state casts doubt upon the interpretation of the charge-exchange experiments. The present results resolve this paradox.

The G-T peak, which is greatly enhanced at forward angles, is much stronger at 160 MeV than at 45 MeV bombarding energy. Its  $1^+$  character has been established on the basis

of excitation energy and the observed angular distributions. In addition to the broad bump observed in the  $^{90}\text{Zr}(p,n)^{90}\text{Nb}$  reaction, about 3.6 MeV above the isobaric analogue of the target ground state (IAS), a second smaller peak is indicated about 8.3 MeV above the IAS<sup>13,19</sup> which also appears to have a predominantly  $\lambda=0$  angular distribution. It has been suggested that this peak is the  $T=5$  ( $T_z$ ) component of the giant G-T resonance. However, it is too weak and too close to the large  $T=4$  ( $T_z$ ) bump to be clearly separated from it.

These results implied that M1 strength might be observable in the parent nucleus using the (p,p') reaction. The expected excitation energy would be given by the difference between the energies of the IAS and the  $T_z$  peak, viz. 8.3 MeV in  $^{90}\text{Zr}$ . The G-T peak in (p,n) reactions is more prominent at higher bombarding energies because the ratio of the spin-flip component,  $V_{\sigma'}$ , to the non-spin-flip component,  $V_{\tau}$ , of the effective interaction increases with energy.<sup>20-24</sup> In the (p,p') reaction above 100 MeV, the same component,  $V_{\sigma'}$ , is mainly responsible for unnatural parity transitions such as  $O^+ \rightarrow I^+$  at small momentum transfer. Since the reaction dynamics should be similar for (p,p') and (p,n) reactions at similar bombarding energies,  $I^+$  states in (p,p') should also be excited mainly through an  $\lambda=0$  transfer. These states should, therefore, show the same

sharp forward peaking of the angular distribution which characterized the G-T peak in the (p,n) reaction.

These considerations prompted a search for M1 strength in medium weight nuclei using the 201 MeV proton beam from the Orsay synchrocyclotron and a large magnetic spectrometer. The targets used were calcium (natural Ca; 15.0 mg/cm<sup>2</sup>),  $^{90}\text{Zr}$  (97.65% enriched; 18.9 mg/cm<sup>2</sup>),  $^{92}\text{Zr}$  (95.13% enriched; 25.4 mg/cm<sup>2</sup>), and  $^{94}\text{Zr}$  (98.6% enriched; 16.4 mg/cm<sup>2</sup>). The particles were detected by two multiwire proportional counters<sup>25</sup> backed by two plastic scintillators. The combination of large size, careful beam preparation, and the trajectory information from the counters allows such clean detection of particles that inelastic scattering data can be taken down to a laboratory angle of 3°. The energy resolution obtained was about 80 keV FWHM. Absolute values of the cross sections were determined by comparison with the known p-p scattering cross sections using a polyethylene target. The spectra were energy calibrated by recording the position of the elastic peak for various magnetic field settings and by using the positions of the known low lying states of  $^{90}\text{Zr}$ ,  $^{40}\text{Ca}$ , and  $^{12}\text{C}$ . The calibration is good to about  $\pm 25$  keV.

Since a known  $I^+$  state at 10.32 MeV has been observed in  $^{40}\text{Ca}$  in a backward angle (e,e') experiment<sup>8</sup>, this nucleus was studied to check that the same state would be selectively

excited in forward angle (p,p') measurements. Many sharp peaks were observed in the  $7^0$  spectrum in the region of excitation energy from 6.8 MeV to above 13 MeV, with the known  $1^+$  state<sup>8</sup> at 10.32 MeV being quite weak. However, at  $3^0$  the spectrum was much simpler, with a peak at 10.3 MeV excitation energy standing out quite clearly above the background. The known  $2^-$  state at 8.43 MeV is also visible in the  $3^0$  proton spectrum. The state at 9.87 MeV is quite strong at  $7^0$  compared to the 10.32 MeV state but is much weaker at  $3^0$ . This is consistent with the previous conclusion that this peak is a close ( $1^+$ ,  $2^+$ ) doublet with the  $1^+$  strength being at most 1/3 that of the 10.32 MeV state.<sup>8</sup>

Spectra from  $^{90}\text{Zr}$ ,  $^{92}\text{Zr}$ , and  $^{94}\text{Zr}$  at a laboratory angle of  $4^0$  are shown in Fig. 1. These spectra show only two features, a broad peak near 9 MeV excitation and a broader peak at about 16 MeV excitation. The centroid of the 16 MeV peak moves toward the elastic peak as the scattering angle increases. This peak is probably a mixture of the E1 giant resonance<sup>26</sup> and the lower lying giant quadrupole and monopole resonances.<sup>27-29</sup> The dipole resonance is expected to peak near  $5^0$  or at even more forward angles due to Coulomb excitation<sup>30</sup>, but the quadrupole resonance should peak near an angle of  $9^0$ . The relative excitation of these three resonances at different angles accounts at least qualitatively for the shift of the centroid with angle.

The peak observed near 9 MeV in all three zirconium isotopes is believed to be the giant M1 resonance. The evidence for this is based on excitation energy, width, angular distribution, and cross section. To ensure that the features observed in the Zr spectra were not an instrumental effect arising from elastic scattering, a spectrum from a gold target was taken at  $4^0$ . No feature could be identified in the gold spectrum even though the elastic scattering count rate at  $4^0$  was much higher than for the zirconium target.

The excitation energy of the peak is consistent both with theoretical predictions<sup>2,4</sup> and with the estimate from the (p,n) data.<sup>13</sup> While this latter value of  $8.3 \pm 0.5$  MeV is rather uncertain, it agrees with the position observed in  $^{90}\text{Zr}$  of  $8.9 \pm 0.2$  MeV. A summary of the excitation energies, widths, and cross sections for the states believed to be  $1^+$  are given in Table I. A recent measurement<sup>19</sup> with fairly good statistics gives a width of 1.8 MeV for the  $T_3$  state in  $^{90}\text{Nb}$ , a value which is consistent with the present (p,p') experiment.

The angular distributions for a number of states in  $^{40}\text{Ca}$ ,  $^{90}\text{Zr}$ , and  $^{94}\text{Zr}$  are shown in Fig. 2. The known  $1^+$  state at 10.3 MeV in  $^{40}\text{Ca}$ , the state at 12.0 MeV in  $^{40}\text{Ca}$ , and the peaks at 8.9 MeV in  $^{90}\text{Zr}$  and 8.6 MeV in  $^{94}\text{Zr}$  all show

the very strong forward peaking which is characteristic of an  $l=0$  angular momentum transfer. This is consistent with the angular distributions observed for the G-T resonances in the (p,n) reaction. If a state observed at 12.0 MeV in  $^{40}\text{Ca}$  is assumed to have natural parity (since it was not excited in the backward angle (e,e') experiment) its spin parity is probably  $0^+$ . The known  $2^-$  state at 8.4 MeV in  $^{40}\text{Ca}$  is expected to be excited dominantly by an  $l=1$  transfer and indeed it does not show a rapid decrease with angle. The broader bump in  $^{90}\text{Zr}$  which probably contains electric dipole, quadrupole, and monopole strengths, also has an angular distribution which is less sharply forward peaked than that expected for the case of a pure  $l=0$ . The fact that the peaks near 9 MeV in  $^{90}\text{Zr}$  and  $^{94}\text{Zr}$  have angular distributions similar to that for the known  $1^+$  state in  $^{40}\text{Ca}$  is strong evidence that these states have spin-parity  $0^+$  or  $1^+$ . However, the fact that no resonance was seen at this excitation energy in  $^{90}\text{Zr}$  ( $\alpha,\alpha'$ )<sup>28,29</sup> implies that the  $J^\pi$  of the resonance is indeed  $1^+$ .

Distorted wave Born approximation (DWBA) calculations have also been carried out using the code DWUCK<sup>31</sup> with macroscopic form factors. The optical parameters have been taken from the systematic studies of Nadasen et al.<sup>32</sup> While this model of form factors is known to be unrealistic for  $l=0$  and  $l$  transfers and cannot be used for absolute comparison

with experiment, these calculations are also known to give angular distribution shapes which are very close to those obtained from more realistic microscopic calculations. For example, the  $l=0$  shape matches very well the experimental angular distribution for the G-T resonance observed in (p,n). As is clear from Fig. 2, these calculations show that the peaks near 9 MeV in  $^{90}\text{Zr}$  and  $^{94}\text{Zr}$  are populated by an  $l=0$  transfer and are therefore consistent with their identification as predominantly M1 transitions.

A further piece of evidence for the M1 nature of the transitions is provided by the strength observed. We assume, as seems consistent with the known values of  $V_{0T}$  and  $V_{0'}$ ,<sup>23,24</sup> that the transition is mediated mainly by  $V_{0T}$ . Following the method given by Goodman et al.,<sup>14</sup> the cross section at zero degrees for  $^{90}\text{Zr}(p,p')$  is estimated to be  $\sim 6$  mb/sr, including a reasonable quenching factor compared to the pure shell model.<sup>33</sup> If we now use the DWBA calculations to extrapolate from  $0^\circ$  to  $4^\circ$ , the predicted cross section for  $^{90}\text{Zr}$  at  $4^\circ$  is  $\sim 3$  mb/sr, which is in good agreement with the 2.8 mb/sr observed.

In summary, resonances about 1.5 to 1.7 MeV wide at excitation energies between 8.6 and 8.9 MeV are observed in the (p,p') reaction on  $^{90}\text{Zr}$ ,  $^{92}\text{Zr}$ , and  $^{94}\text{Zr}$ . A comparison with the angular distribution for a known  $1^+$  state in  $^{40}\text{Ca}$  and with DWBA calculations indicates that these states are

Populated by an angular momentum transfer of zero. The position and cross section of these features suggest that they correspond to M1 transitions observed for the first time in these nuclei.

Acknowledgement:

This work was partly supported by the U.S. National Science Foundation under Grant No. Phy 78-22696. Two of the authors, N.A. and G.M.C., received travel support from the INT Division of NSF.

References:

1. G. Bertsch, L. Zamick, and A. Mekjian, Lecture Notes in Physics, No. 119, Nuclear Spectroscopy, (Springer-Verlag, 1980), p. 245.
2. G.F. Bertsch, Proceedings of the International Conference on Nuclear Physics, Berkeley, Aug. 1980 (to be published).
3. G.E. Brown, J.S. Dehesa, and J. Speth, Nuclear Physics A330, 290 (1979)
4. F.E. Cecil, G.T. Garvey, and W.J. Braithwaite, Nuclear Physics A232, 22 (1974)
5. H. Ikegami et al., Proceedings of 1980 RCNP International Symposium on Highly Excited States in Nuclear Reactions, Osaka, May 1980, p. 250.
6. W. Knüpfner et al., Physics Letters 77B, 367 (1978)
7. A. Richter, Lecture Notes in Physics, No. 108 (Springer, Berlin, 1979) p. 19 and Lecture notes at USSR meeting, 1980 (unpublished).
8. W. Steffan et al., Physics Letters 95B, 23 (1980)
9. G.E. Brown and S. Raman, Comments Nucl. Part. Phys. 9, No. 3, 79 (1980) and D.J. Horen, Proceedings of 1980 RCNP International Symposium on Highly Excited States in Nuclear Reactions, Osaka, May 1980.
10. L.W. Fagg, Rev. Mod. Phys. 47, 683 (1975) and F.E. Cecil et al. quoted in A. Galonsky, The (p,n) Reaction and the Nucleon-Nucleon Force edited by C.D. Goodman et al. (Plenum Press, New York, 1979), p. 191.

Table I:

Characteristics of Resonances Observed with  $l=0$

Target	$E_x$ (MeV)	FWHM (MeV)	$(d\sigma/d\Omega)_{cm}$ ( $4^\circ$ ) (mb/sr)
$^{40}\text{Ca}$	$10.32 \pm 0.02$	$< 0.1$ MeV	$0.24 \pm 0.02$
$^{90}\text{Zr}$	$8.90 \pm 0.15$	$1.7 \pm 0.2$	$2.8 \pm 0.3$
$^{92}\text{Zr}$	$8.8 \pm 0.2$	$1.7 \pm 0.2$	$2.8 \pm 0.3$
$^{94}\text{Zr}$	$8.63 \pm 0.15$	$1.5 \pm 0.2$	$3.1 \pm 0.3$

11. R.R. Doering et al., Phys. Rev. Letters 35, 1691 (1975).
12. A. Galonsky et al., Phys. Letters 74B, 176 (1978).
13. D.E. Bainum et al., Phys. Rev. Letters 44, 1751 (1980).
14. C.D. Goodman et al., Phys. Rev. Letters 44, 1755 (1980).
15. B.D. Anderson et al., Phys. Rev. Letters 45, 699 (1980).
16. D.J. Horen et al., Phys. Letters 95B, 27 (1980).
17. W. Steerrenburg et al., Phys. Rev. Letters 45, 1839 (1980).
18. G. Bertsch, D. Cha, and H. Toki, MSU preprint (1981).
19. W. Steerrenburg et al., Proc. International Conf. Nucl. Phys., Berkeley 1980 and A. Galonsky and T. Nees, private communication.
20. W.G. Love, The (p,n) Reaction and the Nucleon-Nucleon Force, ed. by D.C. Goodman, S.M. Austin, S.D. Bloom, J. Rapaport, and G.R. Satchler, (Plenum Press, New York 1979), p. 23.
21. F. Petrovich, ibid., p. 115.
22. F. Petrovich, W.G. Love, R.J. McCarthy, Phys. Rev. 21C, 1718 (1980).
23. S.M. Austin, The (p,n) Reaction and the Nucleon-Nucleon Force, ed. by C.D. Goodman et al., (Plenum Press, New York 1979), p. 203.
24. W. Steerrenburg et al., Phys. Lett. 91B, 337 (1980).
25. A. Willis et al., Nuclear Physics A144, 137 (1980).
26. B.L. Berman, and S.C. Pultz, Rev. Mod. Phys. 47, 713 (1975)
27. F.E. Bertrand, Ann. Rev. Nucl. Sci. 26, 457 (1976).
28. D.H. Youngblood et al., Phys. Rev. C13, 983 (1976) and D.H. Youngblood, Proceedings of 1980 RCNP International Symposium on Highly Excited States in Nuclear Reactions, Osaka, May 1980, p. 111.
29. M. Buenerd et al., Phys. Let. 84B, 305 (1979) and P. Martin et al., ibid., p. 131.
30. C. Djalali et al., Z. Physik A. 298, 79 (1980).
31. P.D. Kunz, Univ. of Colorado report DWUCK 4 (unpublished).
32. A. Nadasen et al. (to be published)
33. G. Bertsch and H. Toki - private communication.

Figure Captions:

- Fig. 1 Inelastic proton spectra for  $^{90}\text{Zr}$ ,  $^{92}\text{Zr}$ , and  $^{94}\text{Zr}$  at  $4^\circ$ . The arrows indicate the M1 state.
- Fig. 2 Angular distributions for the indicated states in  $^{40}\text{Ca}$ ,  $^{90}\text{Zr}$ , and  $^{94}\text{Zr}$ . The curves are the results of DWBA calculations discussed in the text and are labeled by the  $l$ -transfer of the transitions.



