

GEOMETRIC CHAOTICITY LEADS TO ORDERED SPECTRA FOR RANDOMLY INTERACTING FERMIONS

D. Mulhall, A. Volya and V. Zelevinsky

The interplay of regular and chaotic features in many-body quantum dynamics is currently extensively studied both for simple models and for realistic applications to atomic, nuclear [1], and condensed matter physics, as well as for understanding properties of the QCD vacuum. Typical finite “shell-model” systems, such as complex atoms and nuclei, are described by a mean field and corresponding residual interactions. If there are exact conservation laws the spectra are divided into classes, and one usually deals with the states of each class separately. However, little attention was paid to the problem of correlations between classes of states which are described by the same Hamiltonian but belong to different values of exact integrals of motion.

An obvious and practically important example is angular momentum conservation in a finite Fermi-system. The prediagonalization procedure of projecting the correct value J of nuclear spin out of the m -scheme Slater determinants induces by itself a strong mixing of the states within a shell model configuration [1]. The projected states of various spins acquire a nearly uniform degree of complexity and energy dispersion. For a sufficiently large dimension, the majority of states correspond to a complicated quasi-random coupling of individual spins. This “geometric chaoticity” was used long ago in evaluating the level density for a given J .

A new angle of looking at the problem was introduced by refs. [2, 3] where the spectrum of a random but rotationally invariant Hamiltonian was obtained for a shell-model Fermi system. In spite of the random character of the two-body interaction, the fraction f_0 of the ensemble realizations with a ground state spin $J_0 = 0$ was much higher than the total statistical fraction f_0^s of $J = 0$ states in shell-model space. This result was confirmed as well as for the interacting boson model. A newly-discovered feature was an excess of the probability $f_{J_{max}}$ for the ground state to have the maximum possible spin J_{max} . The emergence of regular features as a result of a random interaction seems to contradict the notion of geometrical chaoticity. Below we show that, vice versa, the geometric chaoticity provides a base for explaining the main features of the pattern.

First we give a couple of trivial examples which point out the possible source of the effects, namely an analog of the Hund rule in atomic physics. Consider a system of N pairwise interacting spins with the Hamiltonian

$$H = A \sum_{a \neq b} \mathbf{s}_a \cdot \mathbf{s}_b = A[\mathbf{S}^2 - Ns(s+1)]. \quad (1)$$

If the interaction strength A is a random variable with zero mean, then the ground state of the system will have equal, $f_0 = f_{S_{max}} = 1/2$, probabilities to have spin $S = 0$ or $S = S_{max}$ (antiferromagnetism or ferromagnetism).

Let us consider a system of interacting fermions. For simplicity we limit ourselves here to a case of N identical particles on a single- j shell which provides a generic framework for the extreme limit of strong residual interaction. Rotational invariance is preserved, so that all single-particle m -states are degenerate in energy. Within this space, the general two-fermion rotationally invariant interaction can be written as $H = \sum_{L\Lambda} V_L P_{L\Lambda}^\dagger P_{L\Lambda}$, where the pair operators with pair spin L and its projection Λ are defined as

$P_{L\Lambda}^\dagger = \frac{1}{\sqrt{2}} \sum_{mn} C_{mn}^{L\Lambda} a_m^\dagger a_n^\dagger$, $P_{L\Lambda} = \frac{1}{\sqrt{2}} \sum_{mn} C_{mn}^{L\Lambda} a_n a_m$ and C are the Clebsch-Gordan coefficients. Because of Fermi statistics, only even L values are allowed in the single- j space. Assuming that the coupling constants V_L are random, uncorrelated and uniformly distributed between -1 and 1, we get the distribution f_j of the ground state spin J_0 shown in Fig. 1(a-e) for $N = 4$ and $N = 6$ at different values of j . For comparison we show by dotted lines the statistical distributions f_j^b based on the fraction of states of given J in the entire Hilbert space for given N . The overwhelming probability f_0 shows the same phenomenon in the uniform ensemble as observed earlier in Gaussian ensembles of V_L [2, 3]. Further evidence of the dominance of $J_0 = 0$ configurations is given by the example, Fig. 1(e), for an odd number of particles, where excess of the ground state spin $J_0 = j$ is evidently related to the ground spin $J_0 = 0$ in the neighboring even system.

First we note that the effect seems to exist already in a crude approximation modeling fermionic pairs by bosons. The commutation relations for the fermion pair operators are (L and L' are even),

$$[P_{L'\Lambda'}, P_{L\Lambda}^\dagger] = \delta_{L'L} \delta_{\Lambda'\Lambda} + 2 \sum_{mm'n} C_{m'n}^{L'\Lambda'} C_{nm}^{L\Lambda} a_m^\dagger a_{m'}. \quad (2)$$

The second term in (2) is of the order N/Ω where Ω is the capacity ($= 2j + 1$ in our case) of the fermionic orbitals. It is small for a small number of fermions; for a nearly filled shell its effect is also small because of the particle-hole symmetry of states. For intermediate shell occupation this term is not small but can be approximately substituted by its mean value (the monopole part with spin $K = 0$). Then, after a simple renormalization, $P_{L\Lambda}$ become bosonic operators, and the Hamiltonian becomes a sum of random bosonic energies $\sum_{L\Lambda} \omega_L n_{L\Lambda}$. The ground state in each realization corresponds to the condensation of the bosons into the single-boson states $|L\Lambda\rangle$ with the lowest value of ω_L . For a given L , the many-boson states with different J allowed for the condensate are degenerate, but the value $L = 0$ is singled out by the obvious fact that for $\omega_0 = \min$ all degenerate states have total spin $J = 0$ while for the minimum boson energy ω_L at $L \neq 0$ any specific value of J , including $J = 0$, appears only in a small fraction of states. If all V_L have the same distribution, we expect $f_0^b \approx 1/k$ where k is a number of (equiprobable) values of L . All other values $J \neq 0$ appear with small probabilities $\sim 1/k^2$. This is demonstrated by Fig. 1(f) where the pattern is qualitatively similar to that in Fig. 1(a-e). The bosonic effect gives only a part (decreasing with increasing j) of the $J_0 = 0$ dominance observed for the fermions. Another argument against the dominance of the bosonic correlations is given in Fig. 1(c). Here we see that after exact elimination of the monopole term ($V_{L=0} \equiv 0$), the picture does not significantly change although the value V_0 is now the lowest only in a small fraction, $\sim 2^{-(k-1)}$, of all cases (when all $V_{L \neq 0}$ are positive).

In our opinion, the main effect comes from the statistical correlations of the fermions. They resolve the bosonic degeneracy in favor of the $J = 0$ and $J = J_{max}$ ground states. In the strong mixing among nearly degenerate states, the eigenstates emerge as complicated chaotic superpositions. The only constraints left are the conservation laws for the particle number and total spin. The latter can be taken into account by the standard cranking approach. Thus, we model the system by the Fermi-gas in statistical equilibrium with the occupation numbers n_m of individual orbitals characterized by the angular momentum projection m onto the cranking axis. The presence of the constraints creates a “body-fixed frame” and splits effective quasiparticle energies, although instead of the collective rotation around a perpendicular axis we have here a random coupling of individual spins with the symmetry (cranking) axis being the only direction




Figure 1: The distribution of ground state angular momenta for various systems of N fermions of spin j , (a–e). The bosonic approximation, f_J^b is in panel (f). The dotted lines are the statistical distribution of allowed J and the solid lines are the ensemble results. In (c) the dashed line is for $V_0 = 0$, i.e. no pairing.

which is singled out in the system. Under the constraints

$$N = \sum_m n_m, \quad M = \sum_m m n_m, \quad (3)$$

equilibrium statistical mechanics leads to the Fermi-Dirac distribution

$$n_m = \frac{1}{\exp(\gamma m - \mu) + 1} \quad (4)$$

determined by the Lagrange multipliers of the chemical potential μ and cranking frequency γ ; in the end the total projection M (equivalent to the K quantum number for axially deformed nuclei) is identified with the total spin J .

The quantities $\mu(N, M)$ and $\gamma(N, M)$ can be found directly from (3). With no cranking, one has the uniform distribution of occupancies $n_m^0 = \bar{n} = N/\Omega$. With the perturbational cranking, the occupation numbers are

$$n_m = \bar{n} \left[1 - \gamma m(1 - \bar{n}) + \frac{\gamma^2}{2}(m^2 - \langle m^2 \rangle)(1 - \bar{n})(1 - 2\bar{n}) + \dots \right]. \quad (5)$$

Here $\langle m^2 \rangle = (1/\Omega) \sum_m m^2 = j^2/3$, and terms of higher orders are not shown explicitly. The expectation value of energy in our statistical system can be written as $\langle H \rangle = \sum_{L\Lambda m_1 m_2} V_L |C_{m_1 m_2}^{L\Lambda}|^2 \langle n_{m_1} n_{m_2} \rangle$. Neglecting the correlations between the occupation numbers, $\langle n_{m_1} n_{m_2} \rangle \approx n_{m_1} n_{m_2}$, we use the statistical result (5) and calculate the geometrical sums with the Clebsch-Gordan coefficients. Expressing the parameter γ in

terms of the total spin $M \rightarrow J$, we come to the result including the terms of the second order in \mathcal{J} ,

$$\langle H \rangle_{N,J} = \sum_L (2L+1) V_L [h_0(L) + h_2(L) J^2 + h_4(L) J^4], \quad (6)$$

where

$$h_0(L) = \bar{n}^2, \quad h_2(L) = \frac{3}{2} \frac{\mathbf{L}^2 - 2\mathbf{j}^2}{\mathbf{j}^4 \Omega^2}, \quad (7)$$

$$h_4(L) = \frac{9}{40} \frac{(1-2\bar{n})^2 (3\mathbf{L}^4 + 3\mathbf{L}^2 - 12\mathbf{j}^2 \mathbf{L}^2 - 6\mathbf{j}^2 + 8\mathbf{j}^4)}{(1-\bar{n})^2 N^2 \Omega^2 \mathbf{j}^8}. \quad (8)$$

J_0 is determined by the ensemble distributions of $h_{2,4} = \sum_L (2L+1) V_L h_{2,4}(L)$. For all realizations of the random interaction with non-negative h_2 and positive h_4 , the ground state has spin $J_0 = 0$. If $h_2 > 0$ but $h_4 < 0$, one has a local minimum of energy at $J = 0$ although there is a possibility to reach the absolute energy minimum at $J_{max} = (1/2)N(\Omega - N)$. This will not happen if at $J = J_{max}$ we still have $h_2 + J_{max}^2 h_4 > 0$. In short, f_0 should be close to 50%, and $f_{J_{max}}$ should be a little less than 50%.

To conclude, we have shown that statistical correlations of fermions in a finite Fermi system with random interactions drive the ground state spin to its minimum or maximum value. This effect is related to the geometrical chaoticity of the random spin coupling of individual particles. This means, that the dominance of 0^+ ground states in even-even nuclei may at least partly come from incoherent interactions rather than solely from coherent pairing. The structure of ground states with an ‘‘antiferromagnetic’’ type ordering, $J_0 = 0$, is compatible with the predictions for chaotic dynamics. Quantitative relations between the effects of geometric chaoticity and pure dynamic effects in finite many-body systems should be an interesting subject for further detailed studies.

The authors wish to acknowledge P. Cejnar whose expertise in the interacting boson model was very helpful. The authors are grateful to G.F. Bertsch, B.A. Brown, V. Cerovski, V.V. Flambaum, M. Horoi, F.M. Izrailev, and D. Kusnezov for constructive discussions. This work was supported by the NSF grant 96-05207.

References

1. V. Zelevinsky *et al.*, Phys. Rep. 276, 85 (1996).
2. C.W. Johnson, G.F. Bertsch and D.J. Dean, Phys. Rev. Lett. 80, 2749 (1998).
3. C.W. Johnson *et al.*, Phys. Rev. C 61, 014311 (2000).